

Crossover behaviors in liquid region of vortex states above a critical point caused by point defects

Yoshihiko Nonomura* and Xiao Hu

Computational Materials Science Center, National Institute for Materials Science, Tsukuba, Ibaraki 305-0047, Japan

(Dated: February 2, 2008)

Vortex states in high- T_c superconductors with point defects are studied by large-scale Monte Carlo simulations of the three-dimensional frustrated XY model. A critical point is observed on the first-order phase boundary between the vortex slush and vortex liquid phases. A step-like anomaly of the specific heat is detected in simulations of finite systems, similar to an experimental observation [F. Bouquet *et al.*, Nature (London) **411**, 448 (2001)]. However, it diminishes with increasing system size, and the number and size distribution of thermally-excited vortex loops show continuous behaviors around this anomaly. Therefore, the present study suggests a crossover rather than a thermodynamic phase transition above the critical point.

PACS numbers: 74.25.Qt, 74.62.Dh, 74.25.Dw

Intensive studies on vortex states of high- T_c superconductors have revealed complex structures of the phase diagram in the magnetic field along the c axis. Point defects such as oxygen deficiencies are essential for such complexity. The Bragg glass (BrG) [1, 2] phase is stable at low temperatures and low magnetic fields. This phase melts into the vortex liquid (VL) or vortex glass (VG) [3] phases as temperature T or magnetic field B increases, respectively. The BrG phase has a closed first-order boundary in the B - T phase diagram. [4, 5] The vortex slush (VS) [6] phase locates between the VG and VL phases, and another first-order boundary between the VS and VL phases terminates at a critical point. [6, 7]

The BrG-VL, [8, 9, 10, 11, 12] BrG-VG [13, 14, 15, 16] and VS-VL [17, 18] phase boundaries have been observed experimentally. Similar phase diagrams have been obtained by numerical simulations including point defects, [19, 20, 21, 22, 23] though the stability of the VG phase is still controversial. [24, 25]

Recently Bouquet *et al.* [26] observed a δ -function peak and a sharp step of the specific heat below and above a critical point, respectively. They argued that this step-like anomaly corresponds to a second-order phase transition related to a vortex-loop blowout. [27, 28]

In the present article, we investigate the phase diagram in the vicinity of the critical point of the VS-VL phase boundary. As displayed in Fig. 1, we find that the critical point is located at $\epsilon \approx 0.15$, which is just above the maximum value of ϵ in our previous work. [23] A step-like anomaly of the specific heat is observed above the critical point in finite systems. However, the size dependence of this anomaly reveals that this step does not indicate a thermodynamic phase transition but a crossover.

In the present study we use the same model as investigated in our previous article on vortex states with point defects, [23] namely the three-dimensional anisotropic,




FIG. 1: Phase diagram of the model (1) in the plane of the strength of point defects (ϵ) and temperature based on a finite system with $L_a = L_b = 50$ and $L_c = 40$. Part of this phase diagram (for $\epsilon \leq 0.14$) has already been published. [23] The first-order VS-VL phase boundary terminates at the critical point (full circle), and the step of the specific heat (dotted line) is observed above the critical point. The VG-VS phase boundary for $\epsilon > 0.14$ is out of the scope in the present study.

frustrated XY model on a simple cubic lattice, [29, 30]

$$\mathcal{H} = - \sum_{i,j \in ab \text{ plane}} J_{ij} \cos(\phi_i - \phi_j - A_{ij}) - \frac{J}{\Gamma^2} \sum_{m,n \parallel c \text{ axis}} \cos(\phi_m - \phi_n), \quad (1)$$

$$A_{ij} = \frac{2\pi}{\Phi_0} \int_i^j \mathbf{A}^{(2)} \cdot d\mathbf{r}^{(2)}, \quad \mathbf{B} = \nabla \times \mathbf{A}, \quad (2)$$

with the periodic boundary condition in all directions. A uniform magnetic field $\mathbf{B}$ is applied along the c axis, and the averaged number of flux lines per plaquette is f . A point defect is introduced as the plaquette surrounded by four weaker couplings and distributed randomly in the ab plane with probability p . Couplings are $J_{ij} = (1 - \epsilon)J$

*Also at Lyman Lab. of Physics, Harvard University, Cambridge, MA 02138; Electronic address: nonomura.yoshihiko@nims.go.jp




FIG. 2: Temperature dependence of (a) the specific heat, (b) the density of dislocations in the ab plane and (c) the ratio of point defects which trap flux lines for $\epsilon = 0.14, 0.15, 0.16, 0.18$ and 0.2 (from bottom to top). The origin of the quantities are varied for each ϵ in (a) and (b).

($0 < \epsilon < 1$) on the point defects, and $J_{ij} = J$ elsewhere. As in our previous article, we fix the parameters $\Gamma = 20$, $f = 1/25$ and $p = 0.003$, and vary the strength of point defects ϵ and temperature T . Configurations of defects are also the same as in our previous work. Monte Carlo simulations are started from a very high temperature, and systems are gradually cooled down. Typical Monte Carlo steps are $4 \sim 14 \times 10^7$ for equilibration, and $2 \sim 16 \times 10^7$ for measurement.

First, we concentrate on the system with $L_a = L_b = 50$ and $L_c = 40$. Temperature dependence of the specific heat C for $0.14 \leq \epsilon \leq 0.2$ is displayed in Fig. 2(a). At $\epsilon = 0.14$, C has a δ -function peak at $T_{sl} = 0.0760 J/k_B$. This peak becomes very small at $\epsilon = 0.15$, and is replaced by a step for $\epsilon \geq 0.16$. This result suggests that the critical point locates at $\epsilon \approx 0.15$. The phase diagram obtained from this finite system is summarized in Fig. 1.

Consistent behaviors are seen in other quantities. Temperature dependence of the density of dislocations in the ab plane ρ_d and the ratio of point defects which trap flux lines r_{tr} are given in Figs. 2(b) and 2(c), respectively. The quantity ρ_d characterizes [7, 23] the VS-VL phase transition. At $\epsilon = 0.14$, ρ_d jumps between $T = 0.072 J/k_B$ and $0.076 J/k_B$, where C sharply increases. This jump becomes much smaller at $\epsilon = 0.15$ and invisible for $\epsilon \geq 0.16$. Keeping possible statistical errors in mind, a tiny hump and dip seem to exist at the temperature region T^* where the specific heat has a step-like anomaly for $\epsilon \geq 0.15$, but the hump and dip are one order smaller than the jump of ρ_d related to the VS-VL phase transition, and therefore considered to be finite-size effects.

At $\epsilon = 0.14$, the quantity r_{tr} gradually increases from $T = 0.070 J/k_B$ as approaching the transition temperature T_{sl} , because the short-range order of flux lines in the VS phase becomes fragile as temperature increases. Then, it rapidly decreases just below T_{sl} , because the mobility of flux lines becomes larger in the coexisting region of the first-order phase transition. This anomaly becomes obscure at $\epsilon = 0.15$ and disappears for $\epsilon \geq 0.16$. Although a tiny hump and dip also seem to exist at T^* for $\epsilon \geq 0.15$, they have an opposite behavior to the cusp at $\epsilon = 0.14$; r_{tr} increases at T^* and drops just above T^* , and therefore have a different origin from the cusp related to the VS-VL phase transition.

Next, in order to explore whether this step-like anomaly of the specific heat is a thermodynamic phase transition or not, its size dependence is investigated. Since flux lines in $\mathbf{B} \parallel \hat{c}$ are one-dimensional objects along the c axis, physical quantities mainly depend on L_c . [31] Even in the vortex liquid region in pure systems, the displacement of flux lines along the ab plane is smaller than the random-walk value $\sim L_c^{1/2}$ due to the blocking by other flux lines. In the present case, this displacement would be even smaller due to the pinning by defects. We thus fix $L_a = L_b = 50$ and change L_c . Temperature dependence of the specific heat for $L_c = 40, 56$ and 80 at $\epsilon = 0.2$ is displayed in Fig. 3(a). As L_c increases, the height of the step of the specific heat becomes smaller. Since it is clearly proportional to $1/L_c$ (Fig. 3(b)), this step is expected to vanish in the thermodynamic limit. The anomaly of the specific heat above the critical point of the VS-VL phase boundary is therefore a crossover rather than a thermodynamic phase transition.

Finally, the number and size distribution of thermally-excited vortex loops are investigated in order to check whether this anomaly of the specific heat is related to the vortex-loop blowout [27, 28] or not. Temperature




FIG. 3: (a) Temperature dependence of the specific heat at $\epsilon = 0.2$ for $L_c = 40, 56$ and 80 . Its origin is varied for each ϵ . (b) c -axis size dependence of the height of the step in (a). The straight line is drawn by the least-squares fitting.

dependence of the number of vortex loops for $0.14 \leq \epsilon \leq 0.2$ is displayed in Fig. 4(a). There exist no vortex loops including pancake vortices at these parameter regions. Size distributions of Josephson loops for $\epsilon = 0.14$ and 0.2 are shown in Figs. 4(b) and 4(c), respectively. [32]

Although the number of vortex loops monotonically increases as temperature increases, no symptom of the vortex-loop blowout is seen around T_{sl} or T^* for each ϵ . This is also the case for the size distribution of Josephson loops. In Figs. 4(b) and 4(c), the number of Josephson loops $N_J(L_J)$ with a given perimeter L_J is plotted versus L_J in a semi-log scale, and the exponential size distribution of $N_J(L_J)$ is clearly observed. There exists no qualitative difference in the behaviors below and above the critical point, and the size distribution changes continuously as temperature is varied around T_{sl} or T^* .

The vortex-loop blowout is characterized by the abrupt increase of the number of vortex loops and the change of the size distribution from an exponential decay to a power-law one at the transition temperature, and the proliferation of vortex loops as large as the system size above the transition temperature. [27, 28] None of these behaviors are observed around T^* in the present simulations, which casts serious doubt on the argument that




FIG. 4: (a) Temperature dependence of the number of vortex loops for $\epsilon = 0.14, 0.15, 0.16, 0.18$ and 0.2 (from bottom to top). Its origin is varied for each ϵ . Size distributions of Josephson loops for (b) $\epsilon = 0.14$ and (c) $\epsilon = 0.2$ around the anomalies of the specific heat are drawn in a semi-log scale.

the step of the specific heat is related to the vortex-loop blowout. [26] In this argument, the vortex-loop blowout scenario was borrowed in order to explain the observation that the value of the specific heat above the step-like anomaly C_+ is larger than that below the anomaly C_- . However, this property is quite general in vortex states, and actually occurs even at the melting transition in pure

systems. [30] In vortex states, the density of vortices increases as temperature increases owing to excitation of vortex loops and wandering of flux lines, and consequently the relation $C_+ > C_-$ holds. The vortex-loop blowout is merely a special case of this general property.

It remains unsolved why a step of the specific heat is clearly observed in experiments even though the sample size is much larger than the system size in our simulations. Several possibilities are: (a) The height of the step of C might saturate at larger L_c , and our maximum system size $L_c = 80$ would be still small. (b) Our model (1) might be still too simple to describe the step of C , e.g. the inhomogeneity of the strength of point defects is not taken into account. (c) The step of C observed experimentally might be a non-equilibrium effect.

In summary, vortex states of high- T_c superconductors with point defects in $\mathbf{B} \parallel \hat{\mathbf{c}}$ has been investigated by large-scale Monte Carlo simulations. The critical point of the first-order phase boundary between the vortex slush and vortex liquid phases is identified as the vanishing point of the δ -function peak of the specific heat. The

shape of the step-like anomaly of the specific heat above the critical point is similar to that of the experiment by Bouquet *et al.* As the system size along the c axis L_c increases, the height of this step seems proportional to $1/L_c$, which suggests that this anomaly is a crossover rather than a thermodynamic phase transition. Consistent behaviors are observed in the temperature dependence of the density of dislocations in the ab plane and the ratio of point defects which trap flux lines. There is no discontinuity in the number and size distribution of vortex loops around the step-like anomaly of the specific heat, which clearly indicates that this anomaly is not related to the vortex-loop blowout.

Numerical calculations were performed on Numerical Materials Simulator (NEC SX-5) at Computational Materials Science Center, National Institute for Materials Science, Japan. Y. N. was supported by the Atomic Energy Research Fund from Ministry of Education, Culture, Sports, Science and Technology, Japan. This study is partially supported by the Priority Grant No. 14038240 from the same Ministry.

-
- [1] T. Nattermann, Phys. Rev. Lett. **64**, 2454 (1990).
 - [2] T. Giamarchi and P. Le Doussal, Phys. Rev. Lett. **72**, 1530 (1994); Phys. Rev. B **52**, 1242 (1995).
 - [3] M. P. A. Fisher, Phys. Rev. Lett. **62**, 1415 (1989); D. S. Fisher, M. P. A. Fisher, and D. A. Huse, Phys. Rev. B **43**, 130 (1991).
 - [4] D. Ertaş and D. R. Nelson, Physica (Amsterdam) **272C**, 79 (1996).
 - [5] T. Giamarchi and P. Le Doussal, Phys. Rev. B **55**, 6577 (1997).
 - [6] T. K. Worthington *et al.*, Phys. Rev. B **46**, 11854 (1992).
 - [7] J. Kierfeld and V. Vinokur, Phys. Rev. B **61**, R14928 (2000).
 - [8] H. Safar *et al.*, Phys. Rev. Lett. **69**, 824 (1992).
 - [9] W. K. Kwok *et al.*, Phys. Rev. Lett. **72**, 1092 (1994).
 - [10] E. Zeldov *et al.*, Nature (London) **375**, 373 (1995).
 - [11] U. Welp *et al.*, Phys. Rev. Lett. **76**, 4809 (1996).
 - [12] A. Schilling *et al.*, Nature (London) **382**, 791 (1996).
 - [13] R. Cubitt *et al.*, Nature (London) **365**, 407 (1993).
 - [14] H. Safar *et al.*, Phys. Rev. B **52**, 6211 (1995).
 - [15] B. Khaykovich *et al.*, Phys. Rev. B **56**, R517 (1997).
 - [16] T. Nishizaki *et al.*, Phys. Rev. B **58**, 11 169 (1998).
 - [17] T. Nishizaki, K. Shibata, and N. Kobayashi, Physica (Amsterdam) **341C-348C**, 957 (2000); K. Shibata *et al.*, Phys. Rev. B **66**, 214518 (2002).
 - [18] H. H. Wen *et al.*, Phys. Rev. B **64**, 054507 (2001).
 - [19] M. J. P. Gingras and D. A. Huse, Phys. Rev. B **53**, 15193 (1996).
 - [20] S. Ryu *et al.*, Phys. Rev. Lett. **77**, 2300 (1996).
 - [21] A. van Otterlo, R. T. Scalettar, and G. T. Zimányi, Phys. Rev. Lett. **81**, 1497 (1998).
 - [22] R. Sugano *et al.*, Physica (Amsterdam) **284B-288B**, 803 (2000), **341C-348C**, 1113 (2000).
 - [23] Y. Nonomura and X. Hu, Phys. Rev. Lett. **86**, 5140 (2001).
 - [24] P. Olsson and S. Teitel, Phys. Rev. Lett. **87**, 137001 (2001).
 - [25] A. Vestergren, J. Lidmar, and M. Wallin, Phys. Rev. Lett. **88**, 117004 (2002).
 - [26] F. Bouquet *et al.*, Nature (London) **411**, 448 (2001).
 - [27] Z. Tešanović, Phys. Rev. B **51**, 16204 (1995), **59**, 6449 (1999).
 - [28] A. K. Nguyen and A. Sudbø, Europhys. Lett. **46**, 780 (1999); Phys. Rev. B **60**, 15307 (1999).
 - [29] Y.-H. Li and S. Teitel, Phys. Rev. Lett. **66**, 3301 (1991); Phys. Rev. B **47**, 359 (1993).
 - [30] X. Hu, S. Miyashita, and M. Tachiki, Phys. Rev. Lett. **79**, 3498 (1997); Phys. Rev. B **58**, 3438 (1998).
 - [31] Y. Nonomura, X. Hu, and M. Tachiki, Phys. Rev. B **59**, R11657 (1999).
 - [32] The effect of intersections of vortex loops is negligible in the present parameter regions, because the averaged number of vortex loops in each ab plane is less than unity, in contrast to the case of Ref. [33].
 - [33] P. Olsson, Europhys. Lett. **58**, 705 (2002).